

BOTX DVD
BOTOX® Cosmetic Examination
TRUE/FALSE
Circle answers

- 1) T / F CME hours are not offered, because trade names and trademarks are involved.
- 2) T / F This is intended for practitioners familiar with general injection techniques & anatomy.
- 3) T / F Cosmetic Botulinum toxin type A is produced and distributed only by Allergan.
- 4) T / F It has FDA approval for glabellar lines; other areas are considered 'off-label' uses.
- 5) T / F All references in this program are to Botulinum toxin type A only.
- 6) T / F The terminal of the nerve is the area of binding and action of Botulinum.
- 7) T / F There is a delay from two days to two weeks for clinical results to be noticed.
- 8) T / F Botulinum toxin is a neurotoxin produced by an aerobic bacterium.
- 9) T / F Like Tetanus, Botulism is a tonic paralysis.
- 10) T / F There are no medical uses for Botulinum toxin.
- 11) T / F Remember to outline 'static' and 'dynamic' wrinkles for your patients.
- 12) T / F Take before and after pictures of your patients.
- 13) T / F Dynamic wrinkles and redundant skin have virtually no response to BOTOX injections.
- 14) T / F Glogau Type one skin has mostly static wrinkles.
- 15) T / F Glogau Type 2 and 3 skin usually responds well to BOTOX injections.
- 16) T / F The side effect of headache-relief may be the reason patients return to this procedure.
- 17) T / F The use of BOTOX Cosmetic is compatible with other types of cosmetic procedures.
- 18) T / F Large temperature fluctuations of the reconstituted medicine should be avoided.
- 19) T / F Dilution should be done with single-use, preservative-free saline.
- 20) T / F One milliliter of saline is required for dilution (i.e. 100 units per cc).
- 21) T / F Men need more Botulinum than women.
- 22) T / F The lateral portion of the Frontalis is a lateral brow elevator.
- 23) T / F The lateral portion of the Obicularis is a lateral brow depressor.

- 24) T / F One should simultaneously paralyze lateral brow depressors & lateral brow elevators.
- 25) T / F The cosmetic portions of the nasal elevators are the superior one-third.
- 26) T / F The use of a topical anesthetic (like the brand-cream, EMLA) is contraindicated.
- 27) T / F Injections are commonly done with 0.5cc, hubless, tuberculin needles (28/30 g, 5/8").
- 28) T / F Insert the needle about 2/3 the length of a 1-1/2 inch needle.
- 29) T / F The glabellar area should be injected with a minimum of 20 units.
- 30) T / F Use 20 to 25 units (total) for the whole forehead.
- 31) T / F For the Frontalis injections, use the superior half of the Frontalis muscle.
- 32) T / F Some patients get relief of tension headaches from cosmetic injections.
- 33) T / F There may be three to six Obicularis injections on either side.
- 34) T / F The middle injection of the Obicularis is lateral to the lateral canthus; one inch from the orbit.
- 35) T / F Do not inject within the rim of the orbit.
- 36) T / F Do not inject medial to the colored part of the eye.
- 37) T / F Do not inject inferior to/below the brow.
- 38) T / F Subcutaneous injections of botulinum are ineffective.
- 39) T / F Facial exercises are an effective way to distribute the medicine after injection.
- 40) T / F The toxin is bound within 4 minutes of injection.
- 41) T / F Know the package insert.
- 42) T / F Do not inject patients taking anticoagulants.
- 43) T / F Injections hurt.
- 44) T / F Aminoglycoside antibiotics may be neurotoxins.
- 45) T / F Treating facial asymmetry is reserved for the experienced cosmetic expert.
- 46) T / F For those with Herpes, Acyclovir prophylaxis is prudent.
- 47) T / F Do not have a "BOTOX Party", if it involves injections.
- 48) T / F Hyperhidrosis responds to local Botulinum injections.
- 49) T / F Contact & involve your state board & malpractice company before using the medicine.

50) T / F Lawyers are not needed for review of your literature and consents.

50 Questions-----EXAMINATION